

The log-canonical threshold of a plane curve

Fernando Hernando

Fernando Hernando (carrilllf@uji.es)
Universidad Jaume I

Carlos Galindo (galindo@mat.uji.ess)
Universidad Jaume I

Francisco Monserrat (framonde@mat.upv.es)
Universidad Politécnica de Valencia

Abstract. We give an explicit formula for the log-canonical threshold of a reduced germ of plane curve. The formula depends only on the first two maximal contact values of the branches and their intersection multiplicities. We also improve the two branches formula given in [4].

References

- [1] E. Artal Bartolo, P. Cassou-Nogués, I. Luengo, A. Melle-Hernández, *On the log-canonical threshold for germs of plane curves*. Contemp. Math. **474** (2008), 1-14.
- [2] N. Budur, P. D. González-Pérez, M. González-Villa, *Log-canonical thresholds of quasi-ordinary hypersurface singularities*. Proc. Amer. Math. Soc. **140** (2012), 4075-4083 .
- [3] T. Järvilehto, *Jumping numbers of a simple complete ideal in a two-dimensional regular local ring*, Ph. D. thesis, University of Helsinki.
- [4] T. Kuwata, *On log canonical thresholds of reducible curves*. Amer. J. Math. **121** (1999), 701-721.
- [5] D. Naie, *Jumping numbers of a unibranch curve on a smooth surface*. Manuscripta Math. **128** (2009), 33-49.
- [6] K. Tucker, *Jumping numbers on algebraic surfaces with rational singularities*. Trans. Amer. Math. Soc. **362** (2010), 3223-3241.